Trabecular Metal[™] **Dental Implant**







As the Trabecular Metal Material interacts with the osteotomy during implant seating, placement will likely have a different tactile feel from other, more conventional implants. The following information is intended to highlight key considerations for placement of the Trabecular Metal Implant to help ensure confidence during surgery.

PREPARATION: Prepare the osteotomy per protocol

Prepare the surgical site by following the Trabecular Metal Implant surgical protocol corresponding to the implant length, diameter and bone density.

DENSE BONE - Do not underprepare the osteotomy in dense bone. If in doubt whether bone type is D2 or D3, dense-bone protocol is recommended. If excess resistance in dense bone is encountered, utilize the appropriate bone tap.

SOFT BONE - Only use the soft-bone protocol in porous trabecular bone with a thin or undifferentiated cortical laver.

PRESSURE: Apply downward pressure to seat

Apply moderate downward pressure as the Trabecular Metal Material begins to engage the osteotomy. The implant may seem to drop into the osteotomy at this point, after which the implant threads re-engage and drive the implant. If the implant needs to be backed out following partial placement into the osteotomy, removal can be achieved by attaching the Surgical Ratchet (RSR) directly to the Fixture Mount/Transfer.

3 TORQUE: Insert implant with low torque values

High insertion torque protocols are not needed for the Trabecular Metal Implant. Placement at 20 Ncm to 35 Ncm is common. Due to the potential for rapid ingrowth and secondary stability, allow the healing process to do its work to achieve stability. If excess resistance in dense bone is encountered, utilize the appropriate bone tap. Osteotomes or any other modifications in the surgical sequence are not recommended.

This guide is for educational use only and should be used as a complement to the standard drilling sequence specified in the Trabecular Metal and Tapered Screw-Vent Implant System Surgical Manual and instructions for use.

Zimmer Biomet Dental **Global Headquarters** 4555 Riverside Drive Palm Beach Gardens, FL 33410 Phone: +1-561-776-6700 Fax: +1-561-776-1272

Contact us at 1-800-342-5454 or visit zimmerbiometdental.com

Unless otherwise indicated, as referenced herein, all trademarks are the property of Zimmer Biomet; and all products are manufactured by one or more of the dental subsidiaries of Zimmer Biomet Holdings, Inc. and marketed and distributed by Zimmer Biomet Dental and its authorized marketing partners. For additional product information, please refer to the individual product labeling or instructions for use. Product clearance and availability may be limited to certain countries/regions. This material is intended for clinicians only and does not comprise medical advice or recommendations. Distribution to any other recipient is prohibited. This material may not be copied or reprinted without the express written consent of Zimmer Biomet Dental. ZBINST0036 REVA 01/19 ©2019 Zimmer Biomet. All rights reserved

